



OPEN
ACCESS

Bioluminescent Mushrooms: Boon for Environmental Health

Waill A. Elkhateeb and Ghoson M. Daba

Chemistry of Natural and Microbial Products Department, National Research Centre, Giza, 12622, Egypt.
Correspondence and requests for materials should be addressed to WAE (email: waillahmed@yahoo.com)

Received:
03-09-2022

Accepted:
10-10-2022

Published:
14-10-2022

Abstract

Mushrooms contain secondary metabolites in their fruit bodies, cultured mycelium, and cultured broth which are higher Basidiomycetes and Ascomycetes. Mushrooms have been used in many sides of human activity for many years. Bioluminescence mushrooms gained more attention. Researchers focused on the ecological role of bioluminescence mushrooms as attracting insects for spore dispersal, which may be beneficial for fungi growing on the forest floor where wind flow is low. In this review, we will focus on some famous bioluminescent mushrooms, their ecology, bioluminescence mechanisms and their ecological benefits.

Keywords: Bioluminescence fungi; Mushroom; *Omphalotus*; *Neonothopanus*; *Armillaria*; *Favolaschia*; *Mycena*

Introduction

Mushrooms are those macrofungi that are known from centuries for their nutritional and medicinal benefits. Thanks to the richness of mushrooms in bioactive compounds that belong to different chemical classes such as phenols, terpenes, proteins, fatty acids, flavonoids, polysaccharides, polyketides, alkaloids, steroids, and other compounds. Ascomycota and Basidiomycota, the two phyla of fungal organisms that include mushrooms, are each known to have both edible and toxic species. Majority of mushrooms are saprophytes, but some are insect parasites, which extend the environmental application of mushrooms as agents that clean environment and recycle essential elements among food chain (ALKolaibe et al., 2021; Daba et al., 2019; Daba et al., 2020; El-Hagrassi et al., 2020; Elkhateeb and Daba, 2020a, b). Also, as clean and efficient biocontrol agents that help in getting rid of many harmful insects and nematodes (Soliman et al., 2022).

On the other hand, many studies have described the activities of the fruiting bodies, crude extracts, and purified compounds originated from mushrooms (Elkhateeb et al., 2019a, b, c, d, e; Elkhateeb and Daba, 2019; Elkhateeb, 2020; Elkhateeb et al., 2020a, b, c, d). These activities include antimicrobial, antiviral, anticancer, antioxidant, antitumor, anti-inflammatory, immunosuppressant, hypocholesterolemic, and hypoglycaemic effects. It should be noted that different species under the same genus may show different biological potentials (Thomas et al., 2019; Thomas et al., 2021; Zhang et al., 2021; Elkhateeb et al., 2021a, b, c, d, e, f). This can be attributed to the change in environmental and growing conditions which are critically affecting the chemical profile of different species. Understanding the importance of mushrooms and their originated compounds encourages for increasing the efforts to discover new species, and study currently identified ones (Elkhateeb and Daba, 2021a, b, c, d, e). There are other mushroom genera that are not edible or even dangerous in addition to the known edible mushroom species. These mushrooms are also generous sources of bioactive compounds that have pharmaceutical and industrial applications. Also, some of these mushrooms can contribute in bio-control of different insects (Elkhateeb et al., 2022a, b; Elkhateeb and Daba, 2022a, b, c, d, e). Natural bioluminescence is the emission of visible lights from a living organism due to a chemical reaction (Haddock et al., 2010). It is widely found across the ocean and terrestrial environments as well as across diverse groups of organisms, including unicellular algae, bacteria (Tanet et al., 2019), multicellular fungi (Karunarathna et al., 2020; Mishra and Srivastava, 2021) and animals (Mensing 2011).



As currently understood, bioluminescence is the consequence of the oxidation of the light-emitting molecule luciferin, which is oxidised by the enzyme luciferase or photoproteins, producing an intermediate high-energy material that then transforms into oxyluciferin (Moline et al., 2013). In this review, we will focus on some famous bioluminescence mushrooms, their ecology, bioluminescence mechanisms of the luminescent fungi, and their benefits.

Bioluminescence phenomenon and Bioluminescent mushrooms

Bioluminescence has been known since ancient times (Weinstein et al., 2016). The phenomenon is most common in marine environments, and a number of theories have been put forward to account for its apparent selective advantage in the dark of the deep ocean (Rees et al., 1998). Hastings (1983) studied the putative ecological functions of bioluminescence and listed defence, offence, and communication; fungal bioluminescence was not covered. A total of 140,000 terrestrial fungus have been found, with about 100 species belonging to four distantly related lineages namely *Armillaria*, *Lucentipis*, *Mycenoid*, and *Omphalotus*, have been recognized as bioluminescent fungi (Kotlobay et al., 2018). Bioluminescent mushrooms emit green light (530 nm) from their fruiting bodies or mycelium when luciferins break down. Oliveira et al. (2012) have found that a single biochemical process is involved across all four bioluminescent monophyletic lineages due to the shared luciferin compound and luciferase enzyme. Numerous studies have suggested that the ecological function of bioluminescence in fungi is to attract insects for spore dispersal, which may be beneficial for fungi growing on the forest floor where wind flow is low (Oliveira et al., 2015).

According to recent experimental research conducted in the Neotropics by Oliveira et al. (2015) for *Neonothopanus gardneri*, it has been hypothesised that the selection advantage offered by bioluminescent fungi (mushroom) can be attributed to the attraction of possibly spore disseminating insects (Agaricales, Marasmiaceae). Alternative explanations might involve warning signals, fungivore deterrence, fungivore predator attraction, or as an unintended byproduct of detoxification or other metabolic processes. Bioluminescence could still have a thermodynamic cost and the ability to attract fungivores. In recent years, the world's known species of bioluminescent fungi (order Agaricales) has dramatically increased. Desjardin et al., (2008), established a list of 64 species of bioluminescent fungi. Chew et al., (2015), revised the list to 81 species. Mihail (2015), discovered five known *Armillaria* species which were bioluminescent. Desjardin et al. (2016), reported two new and two known species of bioluminescent mushrooms from Brazil. Terashima et al. (2016) reported eight new species of bioluminescent mushrooms from Japan. Chew et al. (2011) reported that in Malaysia 71 species of fungi recognised to be luminescent, all Basidiomycetes and belonging to three distinct lineages; *Mycenaceae*, *Omphalotaceae* and *Physalacriaceae*. In China, macrofungi have long been an important part of culture and medicine, and macrofungal diversity in mainland China is becoming more widely studied. However, there have been relatively few reported encounters of bioluminescent mushrooms in the country (Dauner et al., 2021).

Up until recently, a key question has concerned whether molecular oxygen or hydrogen peroxide oxidizes a luciferin substrate in the presence of the enzyme luciferase. Alternatively, it was possible that light emission is a by-product of some metabolic process such as lignin degradation. To create the electrically excited product (oxyluciferin), which decays to the ground state through photon emission, almost all known bioluminescent species utilise oxygen. Thus, the visible and 'cold' light emission results from efficient conversion of energy from chemical bonds to light without heat dissipation (Bechara, 2015). According to a study, the *Neonothopanus gardneri* mushrooms, which are native to the coconut woods of northern Brazil, emit green light under the direction of a circadian clock. Additionally, since light attracts insects, it's possible that bioluminescence aids in the spreading of spores and the spread of fungi (Bechara, 2015). Enchantment and curiosity are immediately suggested when, entering humid woods under a new moon, one sees green, shining mushrooms popping up on the surface of rotten logs. The reader should be able to better appreciate these unusual and beautiful

Agaricomycetes by looking at the picture of a colony of *Mycena lucentipis* mushrooms. Seventy-one out of thousands of fungus species occurring mainly in tropical and temperate zones of the globe are documented to be bioluminescent, of which twenty-six species belong to the genus *Mycena*. As reported in this issue of Current Biology by Oliveira *et al.* (2012), light emission by *Neonothopanus gardneri* mushrooms found in Brazilian coconut forests is controlled by a circadian clock and serves to attract insects for spore dispersal.

Even though the temperate south of the continent is home to various bioluminescent species, including the ubiquitous *Omphalotus nidiformis* (ghost fungus; Agaricales, Marasmiaceae), Weinstein *et al.* (2016) reported that no studies on bioluminescent fungi have focused on the ecological role of bioluminescence. Fungal (Mushroom) bioluminescence uses an enzymatic pathway where a 'luciferin' is oxidised in the presence of 'luciferase' (Desjardin *et al.*, 2008). This process has been tested in several extract-based experiments (Desjardin *et al.*, 2008). In order to show that diverse evolutionary lineages of luminous mushrooms share comparable luciferin-luciferase mechanisms of bioluminescence, Oliveira *et al.* (2012) combined extracts from a variety of species. Basidiomycetes make up all 64 of the mushrooms known to produce bioluminescence (Matheny *et al.*, 2006), but belong to three distinct lineages; the *Omphalotus*, *Armillaria* and the mycenoid lineages (Desjardin *et al.*, 2008). Thus, despite the similarity of the molecular pathways, bioluminescence may have diverse ecological roles in different lineages. According to Weinstein *et al.* (2016), data from several fungus lineages and presumably various growth settings supports both the attraction of spore-dispersing insects and the incidental by-product of metabolism theories for bioluminescence in mushrooms. One possible explanation for such a situation would be that bioluminescence in fungi arose as an incidental by-product of metabolism, and that some fungi such as *Omphalotus nidiformis* remained in this condition. Others, like *Neonothopanus gardneri*, gained a fitness advantage by modifying their bioluminescence to increase their appeal to potential spore-dispersing insects.

***Favolaschia* (Pat.)**

Favolaschia (Pat.) is a saprobic, often tiny, mushroom-like genus of the Basidiomycetes family Mycenaceae, which has a wide range of species. It was reported from a lowland warm-temperate to subtropical and tropical zone (Fig. 1). The genus *Favolaschia* comprises 109 species (Nimalrathna *et al.*, 2022). Hymenial pores, a gelatinous trama, and the presence of gloeocystidia and acanthocystida are characteristics of this genus (Gillen *et al.*, 2012). To date, *F. manipularis* (Audrey *et al.*, 2015) and *F. peziziformis* (Bodensteiner *et al.*, 2004) have been reported to have bioluminescence.




Fig. 1. *Favolaschia* mushroom showing bioluminescent in the dark.

Cited in: <https://imgur.com/gallery/KPzbH>

Omphalotus nidiformis

Omphalotus nidiformis is an Australian basidiomycete from the temperate zone that is typically found around or close to the foot of eucalyptus trees. Its basidomes range in colour from cream to brown and are typically between 20 and 40 cm in diameter (Weinstein *et al.*, 2016). In the

Southern Hemisphere winter (June to July), it develops basidomes and continuously gives off a flimsy white-blue glow (Fig. 2).



Fig. 2. *Omphalotus nidiformis* showing bioluminescent in the dark

Cited in: <https://texasmycology.com/shop/ols/products/mushroom-fruiting-kit-beech-hypsizygu-tesselatus>

Neonothopanus gardneri

Neonothopanus gardneri is a species of bioluminescent fungus belonging to the order Agaricales (Marasmiaceae) found in South America. Its existence was first reported in 1840 by George Gardner in his travels to Brazil, where it is popularly called "coco flower" (Weinstein et al., 2016). Recently, *N. gardneri* had some of its bioactives separated and their corresponding structures clarified. It is mostly found in decomposing leaves and in the trunk of dwarf palm trees such as "pindoba" (*Attalea oleifera*) or "babaçu" (*Orbignya phalerata*) (Fig. 3).



Fig. 3. *Neonothopanus gardneri* showing bioluminescent in the dark

Cited in: <https://www.sciencenews.org/article/how-mushroom-gets-its-glow>

Mycena lucentipis

Mycena is a large genus of small saprotrophic mushrooms that are rarely more than a few centimeters in width. They are characterized by a white spore print, a small conical or bell-shaped cap, and a thin fragile stem. Most are gray or brown, but a few species have brighter colors. Most have a translucent and striate cap, which rarely has an incurved margin. The gills are attached and usually have cystidia (Fig. 4).

According to reports, seven species of *Mycena*, representing specimens gathered in Belize, Brazil, the Dominican Republic, Jamaica, Japan, Malaysia, and Puerto Rico are luminous. Four of them represent new species (*Mycena luxaeterna*, *M. luxarboricola*, *M. luxperpetua* and *M. silvaelucens*) and three represent new reports of luminescence in previously described species (*M. aff. abieticola*, *M. aspratilis* and *M. margarita*). The genus *Mycena* includes about 500 known

species of gilled mushrooms. More than 58 species are known to be bioluminescent (Desjardin et al., 2010).




Fig. 4. *Mycena* sp. showing bioluminescent

Cited in: <https://www.flickr.com/photos/stephenmudge/39774574525>

Armillaria

Armillaria is one of the most important genera of fungal root pathogens worldwide. It attacks hundreds of tree species in both timber (*Abies*, *Picea*, *Pinus* and *Pseudotsuga*). *Armillaria* genus belonging to Basidiomycota, (Physalacriaceae) (Baumgartner et al., 2011). The honey fungus species *Armillaria mellea*, which inhabits trees and woody bushes, belongs to the genus *Armillaria* of fungi. It includes about 10 species formerly categorized summarily as *Armillaria mellea*. The largest known species is *Armillaria ostoyae*. Some species of *Armillaria* display bioluminescence, resulting in foxfire. *Armillaria* can be a destructive forest pathogen. It causes "white rot" root disease of forests. *Armillaria* can kill its host because it is a facultative saprophyte that also consumes dead plant matter, as opposed to parasites that must control their growth to prevent host death (Kirk et al., 2008). The properties and mechanisms of bioluminescence in fungi are also of interest to mycological research (Fig. 5).



Fig. 5. *Armillaria* showing bioluminescent in the dark

Cited in: <https://www.sciencelearn.org.nz/images/3717-armillaria-novae-zelandiae-te-harore>

Unlike the blue light of marine organisms and the yellow light of fireflies, bioluminescent fungi emit greenish-bluish light at a wavelength of 520–530 nm (Ilundu & Okiti 2016). In *Omphalotus* and *Mycenaceae* fungi, luminescence often occurs in both the fruiting bodies and the mycelia, but *Armillaria*, curiously, exhibit luminescence only in the mycelia, producing the "foxfire" effect of glowing wood that has been described for millennia (Desjardin et al., 2008). The biochemical mechanism of bioluminescence in *Armillaria*, *Favolaschia*, *Omphalotus*, *Neonothopanus* and *Mycena* (Basidiomycetes) remains unclear but most research theories proposed both enzymatic and non-enzymatic reaction (Baumgartner et al., 2011).

Benefits and application for bioluminescent fungi

Many different applications for bioluminescent fungi have been formulated, including using them as an indicator for low concentrations of environmental pollutants (Wan et al., 2014) and expanding the use of luminescent markers in biomedicine and biotechnology (Ilundu and Okiti, 2016).

Conclusion

Bioluminescence has been known since ancient times from different living organisms (Marine Fish, plant, bacteria, fungi and others). Bioluminescence fungi (Mushroom) gained more attention from the researchers. Many scientific researches focused on the ecological role of bioluminescence and their benefits. The most famous bioluminescence mushroom were *Armillaria*, *Favolaschia*, *Omphalotus*, *Neonothopanus* and *Mycena* all belonging to Basidiomycetes. Now many researchers discovered many new different mushroom genera. Further studies are required to explore the additional applications of this phenomenon.

References

- ALKolaibe AG, Elkhateeb WA, Elnahas MO, et al. (2021) Wound Healing, Anti-pancreatic Cancer, and α -amylase Inhibitory Potentials of the Edible Mushroom, *Metacordyceps neogunnii*. Research Journal of Pharmacy and Technology 14(10): 5249–5253.
- Audrey L, Desjardin D, Tan Y, et al. (2015) Bioluminescent fungi from Peninsular Malaysia—a taxonomic and phylogenetic overview. Fungal Diversity 70(1): 149–187.
- Baumgartner K, Coetzee M and Hoffmeister D (2011) Secrets of the subterranean pathosystem of *Armillaria*. Molecular Plant Pathology 12(6): 515–534.
- Bechara, EJ (2015) Bioluminescence: A fungal nightlight with an internal timer. Current Biology 25(7): R283–R285.
- Bodensteiner P, Binder M, Moncalvo J, et al. (2004) Phylogenetic relationships of cyphelloid homobasidiomycetes. Molecular Phylogenetics and Evolution 33: 501–515.
- Chew A, Desjardin D, Tan Y, et al. (2015) Bioluminescent fungi from Peninsular Malaysia—a taxonomic and phylogenetic overview. Fungal diversity 70(1): 149–187.
- Chew A, Tan Y, Musa M, et al. (2011) Biodiversity and biology of bioluminescent fungi. Malaysian Society for Microbiology. 460–464.
- Daba GM, Elkhateeb W, ELDien AN, et al. (2020) Therapeutic potentials of n-hexane extracts of the three medicinal mushrooms regarding their anti-colon cancer, antioxidant, and hypocholesterolemic capabilities. Biodiversitas Journal of Biological Diversity 21(6): 1–10.
- Daba GM, Elkhateeb WA, Wen TC, et al. (2019) The continuous story of truffle plant interaction. pp. 375–383. In: Kumar V, Prasad R, Kumar M, Choudhary, D.K. (eds.) Microbiome in Plant Health and Disease, Springer, Singapore.
- Dauner L, Karunarathna S, Tibpromma S, et al. (2021) Bioluminescent fungus *Roridomyces viridiluminus* sp. nov. and the first Chinese record of the genus *Roridomyces*, from South-western China. Phytotaxa 487(3): 233–250.
- Desjardin DE, Oliveira AG and Stevani CV (2008) Fungi bioluminescence revisited. Photochemical and Photobiological Science 7: 170–182.
- Desjardin D, Perry B, Lodge D, et al. (2010) Luminescent *Mycena*: new and noteworthy species. Mycologia 102(2): 459–477.
- Desjardin D, Perry B and Stevani C (2016) New luminescent mycenoid fungi (Basidiomycota, Agaricales) from São Paulo State, Brazil. Mycologia 108(6): 1165–1174.

- El-Hagrassi A, Daba G, Elkhateeb W, et al. (2020) In vitro bioactive potential and chemical analysis of the n-hexane extract of the medicinal mushroom, *Cordyceps militaris*. *Malays J Microbiol* 16(1): 40–48.
- Elkhateeb W, Elnahas M and Daba G (2021e) Infrequent Current and Potential Applications of Mushrooms, CRC Press, 70–81.
- Elkhateeb W, Elnahas MO, Paul W, et al. (2020b) *Fomes fomentarius* and *Polyporus squamosus* models of marvel medicinal mushrooms. *Biomed Res Rev.* 3: 119.
- Elkhateeb W, Thomas P, Elnahas M, Daba G (2021f) Hypogeous and Epigeous Mushrooms in Human Health. *Advances in Macrofungi* 7–19.
- Elkhateeb WA and Daba G (2020) The endless nutritional and pharmaceutical benefits of the Himalayan gold, *Cordyceps*; Current knowledge and prospective potentials. *Biofarmasi Journal of Natural Product Biochemistry* 18(2): 70–77.
- Elkhateeb WA, Daba GM, El-Dein AN, et al. (2020c) Insights into the in-vitro hypocholesterolemic, antioxidant, antirotavirus, and anticolon cancer activities of the methanolic extracts of a Japanese lichen, *Candelariella vitellina*, and a Japanese mushroom, *Ganoderma applanatum*. *Egyptian Pharmaceutical Journal* 19(1): 67.
- Elkhateeb WA, Daba GM, Elmahdy EM, et al. (2019b) Antiviral potential of mushrooms in the light of their biological active compounds. *ARC J Pharmac Sci* 5: 8–12.
- Elkhateeb WA, Daba GM, Elnahas M, et al. (2020d) Metabolic profile and skin-related bioactivities of *Ceriporus squamosus* hydromethanolic extract. *Biodiversitas J Biological Div* 21(10): 4732-4740.
- Elkhateeb WA, Daba GM, Elnahas MO, et al. (2019c) Anticoagulant capacities of some medicinal mushrooms. *ARC J Pharma Sci* 5: 12–16.
- Elkhateeb WA, Daba GM and Gaziea SM (2021d) The Anti-Nemic Potential of Mushroom against Plant-Parasitic Nematodes. *Open Access Journal of Microbiology & Biotechnology* 6(1): 1–6.
- Elkhateeb WA, Daba GM, Sheir D, et al. (2019e) GC-MS analysis and in-vitro hypocholesterolemic, anti-rotavirus, anti-human colon carcinoma activities of the crude extract of a Japanese *Ganoderma* spp. *Egyptian Pharmaceutical Journal* 18(2): 102-110.
- Elkhateeb WA, Daba GM, Thomas PW, et al. (2019d) Medicinal mushrooms as a new source of natural therapeutic bioactive compounds. *Egypt Pharmaceu J* 18(2): 88–101.
- Elkhateeb WA and Daba GM (2019) The amazing potential of fungi in human life. *ARC J Pharma Sci* 5(3): 12–16.
- Elkhateeb WA and Daba GM (2020) *Termitomyces* Marvel Medicinal Mushroom Having a Unique Life Cycle. *Open Access Journal of Pharmaceutical Research* 4(1): 1–4.
- Elkhateeb WA and Daba GM (2021a) Highlights on the Golden Mushroom *Cantharellus cibarius* and unique Shaggy ink cap Mushroom *Coprinus comatus* and Smoky Bracket Mushroom *Bjerkandera adusta* Ecology and Biological Activities. *Open Access Journal of Mycology & Mycological Sciences* 4(2): 1–8.
- Elkhateeb WA and Daba GM (2021b) Highlights on the Wood Blue-Leg Mushroom *Clitocybe Nuda* and Blue-Milk Mushroom *Lactarius Indigo* Ecology and Biological Activities. *Open Access Journal of Pharmaceutical Research* 5(3): 1–6.
- Elkhateeb WA and Daba GM (2021c) Highlights on Unique Orange Pore Cap Mushroom *Favolaschia* Sp. and Beech Orange Mushroom *Cyttaria* sp. and Their Biological Activities. *Open Access Journal of Pharmaceutical Research* 5(3): 1–6.

- Elkhateeb WA and Daba GM (2021d) Mycotherapy of the good and the tasty medicinal mushrooms *Lentinus*, *Pleurotus*, and *Tremella*. *Journal of Pharmaceutics and Pharmacology Research* 4(3): 1–6.
- Elkhateeb WA and Daba GM (2021e) The Fascinating Bird's Nest Mushroom, Secondary Metabolites and Biological Activities. *International Journal of Pharma Research and Health Sciences* 9 (1): 3265–3269.
- Elkhateeb WA and Daba GM (2022a) Medicinal mushroom what should we know. *International Journal of Pharmaceutical Chemistry and Analysis* 9(1): 1-19.
- Elkhateeb WA and Daba GM (2022b) The wild non edible mushrooms, what should we know so far?. *International Journal of Advanced Biochemistry Research* 6(1): 43-50.
- Elkhateeb WA and Daba GM (2022c) Bioactive Potential of Some Fascinating Edible Mushrooms *Flammulina*, *Lyophyllum*, *Agaricus*, *Boletus*, *Letinula*, and *Pleurotus* as a Treasure of Multipurpose Therapeutic Natural Product. *Pharm Res* 6(1): 1-10.
- Elkhateeb WA and Daba GM (2022d) Bioactive Potential of some Fascinating Edible Mushrooms *Macrolepiota*, *Russula*, *Amanita*, *Vovariella* and *Grifola* as a Treasure of Multipurpose Therapeutic Natural Product. *J Mycol Mycological Sci* 5(1): 1-8.
- Elkhateeb WA, El Ghwas DE, Gundoju NR, et al. (2021a) Chicken of the Woods *Laetiporus Sulphureus* and *Schizophyllum Commune* Treasure of Medicinal Mushrooms. *Open Access Journal of Microbiology & Biotechnology* 6(3): 1–7.
- Elkhateeb WA, El-Ghwas DE, Daba GM (2021b) A Review on Ganoderic Acid, Cordycepin and Usnic Acid, an interesting Natural Compounds from Mushrooms and Lichens 5(4): 1–9.
- Elkhateeb WA, EL-Ghwas DE, Daba GM (2022) Mushrooms as Efficient Enzymatic Machinery. *J Biomed Res Environ Sci* 3(4): 423-428.
- Elkhateeb WA, Elnahas M, Wenhua L, et al. (2021c) The coral mushrooms *Ramaria* and *Clavaria*. *Studies in Fungi* 6(1): 495–506.
- Elkhateeb WA, Elnahas MO, Thomas PW, et al. (2019a) To Heal or Not to Heal? Medicinal Mushrooms Wound Healing Capacities. *ARC Journal of Pharmaceutical Sciences* 5(4): 28–35.
- Elkhateeb WA, Elnahas MO, Thomas PW, et al. (2020a) *Trametes Versicolor* and *Dictyophora Indusiata* Champions of Medicinal Mushrooms. *Open Access Journal of Pharmaceutical Research* 4(1): 1–7.
- Elkhateeb WA, Karunarathna SC, Galappaththi MCA, et al. (2022) Mushroom biodegradation and their role in mycoremediation. *Studies in Fungi* (Under press)
- Elkhateeb WA. (2020) What medicinal mushroom can do? *Chem Res J* 5(1): 106–118.
- Gillen K, Læssøe T, Kirschner R, et al. (2012) *Favolaschia* species (Agaricales, Basidiomycota) from Ecuador and Panama. *Nova Hedwigia* 96 (1–2): 117–165.
- Haddock SH, Moline MA, Case JF (2010) Bioluminescence in the sea. *Annual review of marine science* 2: 443–493.
- Hasting J (1983) Biological diversity, chemical mechanisms, and the evolutionary origins of bioluminescent systems. *Journal of Molecular Evolution* 19: 309–321.
- Ilondu E and Okiti A (2016) Bioluminescence in mushroom and its application potentials. *Nigerian Journal of Science and Environment* 14(1): 1-10.
- Karunarathna SC, Mortimer PE, Tibpromma S, et al. (2020) *Roridomyces phyllostachydis* (Agaricales, Mycenaceae), a new bioluminescent fungus from Northeast India. *Phytotaxa* 459(2): 155–167.

- Kirk P, Cannon P, Minter D, Stalpers J (2008) Dictionary of the Fungi Wallingford. UK: CABI, 335.
- Kotlobay AA, Sarkisyan KS, Mokrushina YA, et al. (2018) Genetically encodable bioluminescent system from fungi. Proceedings of the National Academy of Sciences 115 (50): 12728–12732.
- Life. Open Access Journal of Mycology & Mycological Sciences, 5(1): 1–5.
- Matheny PB, Curtis J M, Hofstetter V, et al. (2006) Major clades of Agaricales: a multilocus phylogenetic overview. Mycologia 98: 982–995.
- Mensing AF (2011) Bioluminescence in Fishes. Encyclopaedia of Fish Physiology. Elsevier, USA 497–503.
- Mihail JD (2015) Bioluminescence patterns among North America *Armillaria* species. Fungal Biol 119(6): 528–537.
- Mishra M and Srivastava D (2021) Bioluminescent fungi: reviewing nature's riddle!. Journal of Mycopathology Research. 59(3): 199–206.
- Moline MA, Oliver MJ, Orrico C, et al. (2013) Bioluminescence in the sea. In: Subsea Optics and Imaging. Elsevier 134–170.
- Nimalrathna, TS, Tibpromma, S, Nakamura, A, et al. (2022) The case of the missing mushroom: a novel bioluminescent species discovered within *Favolaschia* in south-western China. Phytotaxa 539(3): 244–256.
- Oliveira AG, Desjardin DE, Perry BA, et al. (2012) Evidence that a single bioluminescent system is shared by all known bioluminescent fungal lineages. Photochemical and Photobiological Sciences 11: 848–852.
- Oliviera AG, Stevani CV, Waldenmaier HE, et al. (2015) Circadian control sheds light on fungal bioluminescence. Current Biology 25: 964–968.
- Rees J-F, de Wergifosse B, Noiset O, et al. (1998) The origins of marine bioluminescence: turning oxygen defence mechanisms into deep-sea communication tools. Journal of Experimental Biology 201: 1211–1221.
- Soliman G, Elkhateeb WA, Wen TC, Daba G (2022) Mushrooms as efficient biocontrol agents against the root-knot. Egyptian Pharmaceutical Journal 1687:4315.
- Tanet L, Tamburini C, Baumas C, et al. (2019) Bacterial bioluminescence: light emission in photobacterium phosphoreum is not under quorum-sensing control. Frontiers in Microbiology 10: 1–9.
- Terashima Y, Takahashi H and Taneyama Y (2016) The fungal flora in south-western Japan: Agarics and boletes. Tokai Daigaku, Kanagawa, Japan 349.
- Thomas P, Elkhateeb W and Daba G (2021) Industrial Applications of Truffles and Truffle-like Fungi. In Advances in Macrofungi 82-88.
- Thomas PW, Elkhateeb WA and Daba G (2019) Truffle and truffle-like fungi from continental Africa. Acta Mycol 54(2): 1–15.
- Wan N, Jusoh W and Wong L (2014) Explaining the potential of whole cell biosensor: A review in environmental applications. International Journal of Chemical, Environmental and Biological Sciences 2 (1): 52–56.
- Weinstein P, Delean S, Wood T, et al. (2016) Bioluminescence in the ghost fungus *Omphalotus nidiformis* does not attract potential spore dispersing insects. IMA fungus 7(2): 229–234.
- Zhang X, Zhang X, Gu S, et al. (2021) Structure analysis and antioxidant activity of polysaccharide-iron (III) from *Cordyceps militaris* mycelia. International Journal of Biological Macromolecules 178: 170-179.

Author Contributions

WAE and GMD conceived the concept, wrote and approved the manuscript.

Acknowledgements

Not applicable.

Funding

There is no funding source for the present study.

Availability of data and materials

Not applicable.

Competing interest

The authors declare no competing interests.

Ethics approval

Not applicable.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution, and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third-party material in this article are included in the article's Creative Commons license unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. Visit for more details <http://creativecommons.org/licenses/by/4.0/>.

Citation: Elkhateeb WA and Daba GM (2022) Bioluminescent Mushrooms: Boon for Environmental Health. Environ Sci Arch 1(2): 88-97.